

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**




Fig. 1




Fig. 2




Fig 1




Fig 2




Fig. 3




Fig 4




Fig 5




Fig 7

Differentiating Acute Myocardial Infarction
From Other ECG Abnormalities

Alan Andresen, Ronald Selvester M.D.

Inovise Medical, Inc.

Clinical Relevance

(Incidence of confounders, difficulty in diagnosis of acute MI)

- LBBB ~5%; Vpacing ~9%
 - Very confusing for overreading clinician; often ignored
- RBBB ~6%
 - Guidelines not adjusted for this confounder (esp. anterior MI)
- LVH ~20% (by conventional ECG criteria)
 - LVH changes on ECG confounder for both sensitivity and specificity (esp. anterior MI)
- Inferior MI w/RV involvement
 - ~40% of RCA occlusions involving RV, with 8x relative risk
 - Clinician identification highly variable; absent for algorithms
- Non-STE ~60% of AMI
 - Borderline STE in this population ~2x for females as compared to males

ECG Algorithm Confounder Performance

Long Beach AMI. n=1151, CK-MB \geq 9 for rule-in

Method	LBBB n=52	RBBB n=90	LVH n=114	RVH n=27	V-pacing n=55	STE ⁻ Female n=263	RV ext n=64
Sensitivity	15%	56%	46%	52%	~15%	29%	~80%
CV	2%	31%	11%	30%	-	7%	-
Alg 1	-	29%	25%	19%	-	10%	-
Alg 2	-	29%	25%	48%	-	17%	-
Alg 3	-	29%	25%	48%	-	17%	-

ECG Algorithm Confounder Performance

MCV AMI, n=1274, CKMB/Troponin+ for rule-in

Method	LB	BB	RVH	V-pacing	STE- Female	RV ext
Sensitivity	n=26	n=39	n=265	-	n=405	-
CV	12%	38%	26%	-	15%	-
Alg 1	4%	31%	6%	-	7%	-
Alg 2	-	-	-	-	-	-
Alg 3	-	10%	19%	-	9%	-

AMI Criteria When RBBB+

- Adjust ST deviation to remove portion of STT abnormality due to block
 - ST baseline adjusted by 10% of difference between max and min amplitudes of final 1/3 of QRS complex
- Suppress detection of AMIs that are unseparable from RBBB
 - Posterior (-V1 to -V3, V6)

AMI Criteria When LBBB+

- Adjust ST deviation to remove portion of STT abnormality due to block
 - ST baseline adjusted by 10% of difference between (Rmax-STJ20) and (Smax-STJ20)
- Adjust STE deviation threshold for calling acute
 - Limb leads: increase by 50 μ V (100 to 150)
 - Precordial leads: increase by 75 μ V (200 to 275)
- Suppress detection of acute MIs that are unseparable from LBBB
 - Anterior (V1-V5)
 - Subendocardial injury
- Add qualifiers to improve specificity
 - Posterior AMI only when non-adjusted depression present in V1-V3

AMI Criteria With ECG Evidence for LVH

- Adjust ST deviation to remove portion of STT abnormality due to "LVH with Repol"
 - ST baseline adjusted by 5% of difference between (Rmax-STJ20) and (Smax-STJ20)
- Adjust ST deviation to remove portion of STT abnormality due to "LVH without Repol"
 - Limb: ST baseline adjusted by 2.5% of difference between (Rmax-STJ20) and (Smax-STJ20)
 - Precordial: ST baseline adjust by lead by -50 to +50uV

Questions for Discussion

- Pursuing AMI detection when confounders present
 - Is there a consensus about clinical relevance?
 - Is it possible to distinguish these confounders from AMI?
 - Are females significantly under-diagnosed for STE AMI?
- If yes to the above...
 - Which is the most important, clinically?
 - Which is the most likely to yield to investigation?
- If no...
 - Where should we focus our AMI efforts?